

Representations of Codimension ≥ 3 by Definite Quadratic Forms

Wai Kiu Chan and Dennis R. Estes

Department of Mathematics, University of Southern California, Los Angeles, California 90089

and

M. Jöchner

Franz-List-Weg 11, 70794 Harthausen, Germany

Communicated by J. S. Hsia

[View metadata, citation and similar papers at core.ac.uk](#)

Let M be a positive definite quadratic \mathbf{Z} -lattice of rank $\geq n + 3$. If N is a quadratic \mathbf{Z} -lattice of rank n which is primitively represented by the genus of M and if all the successive minima of N increase sufficiently quickly, then there exists a global primitive representation of N by M with approximation and primitivity properties.

© 1998 Academic Press

1. INTRODUCTION

Let M be a positive definite quadratic \mathbf{Z} -lattice of rank m , and N a quadratic \mathbf{Z} -lattice of rank n which is primitively represented by M locally. Theorem 2.5 in [3] shows that there exists a global representation of N by M with approximation and primitivity properties provided $m \geq 3n + 3$ and the minimum of N is sufficiently large. Recent work by Kitaoka and Jöchner [4, 5] suggests that $m \geq 2n + 3$ may suffice. This is the case when $n = 1$ (see the remark at the end of [5]) and $n = 2$ [4]. When $n = 1$, one can even push down to $m \geq 4$, see Theorem 2.1 in [2]. In this article, we generalize the result to the case where n is arbitrary and $m \geq n + 3$ by imposing conditions on the sizes of the successive minima of N . For $i = 1, \dots, n$, let $S_i(M, n)$ be the set of positive rational numbers which are the i th successive minima of rank n \mathbf{Z} -lattices primitively represented by the genus of M . The following is our main theorem.

THEOREM 1.1. *Let n be a positive integer and M a positive definite quadratic \mathbf{Z} -lattice of rank $\geq n + 3$. Let s be a positive integer and let T be a*

finite set of primes. There exist real valued functions $c_1 = c_1(M, T, s)$, $c_2 = c_2(M, T, s, \lambda_1)$, ..., $c_n = c_n(M, T, s, \lambda_{n-1})$, $\lambda_i \in S_i(M, n)$, with the following property:

Let N be a positive definite rank n \mathbf{Z} -lattice with successive minima $\mu_1 \leq \dots \mu_n$ and $f_p: N_p \rightarrow M$ primitive representations at each prime p . If $\mu_1 \geq c_i(M, T, s, \mu_{i-1})$ for $i=1, \dots, n$, then there exists a global representation $f: N \rightarrow M$ satisfying the conditions

1. $f \equiv f_p \pmod{p^s M_p}$ for all $p \in T$,
2. $f(N)$ is primitive in M ,

We refer the reader to [1] and [7] for any undefined terminology and notation. In particular, $s(M)$ denotes the scale of M . All lattices and the associated quadratic forms are understood to be positive definite. Given a nonzero vector w in the ambient space V of M , let $\Phi_w: V \rightarrow V$ be the projection $\Phi_w(x) = x - ((w, x)/(w, w))w$ onto the orthogonal complement of w , where $(,)$ is the bilinear form on M . For any $w \in M$, we define $\gamma_w = Q(w)/s(M)$. Then γ_w is an ideal of \mathbf{Z} . By abuse of notation, we also use γ_w to denote a generator of this ideal. Note that $\gamma_w \Phi_w(N) \subseteq N$ for any submodule N of M containing w . Linearly independent vectors $v_1, \dots, v_n \in V$ are Hermite reduced if the corresponding quadratic form $f(x_1, \dots, x_n) = Q(x_1 v_1 + \dots + x_n v_n)$ is Hermite reduced. We refer the reader to Chapter 12 of [1] for the facts needed below for Hermite reduced forms and Siegel domains. In particular, Hermite reduced is defined inductively by $Q(v_1)$ is the minimum nonzero norm on $N = \mathbf{Z}[v_1, \dots, v_1]$, $|2(v_1, v_i)| \leq Q(v_1)$, and $\Phi_{v_1}(v_2), \dots, \Phi_{v_1}(v_n)$ is Hermite reduced. Any vector in N of minimal nonzero norm can be extended to a Hermite reduced basis for N and the form $f(x_1, \dots, x_n) = Q(x_1 v_1 + \dots + x_n v_n)$ for a Hermite reduced basis lies in the Siegel domain $\mathcal{S}_n(4/3, 1/2)$.

LEMMA 1.1. *Let v be a vector in N of minimal nonzero norm. Let μ_1, \dots, μ_n denote the successive minima for N and $\mu'_1, \dots, \mu'_{n-1}$ the successive minima for $N' = \Phi_v(N)$. Then there is a positive constant $C = C(n)$ such that $(\mu_{i+1})/\mu_i, (\mu'_i)/\mu_{i+1} \leq C$ for $i = 1, \dots, n-1$.*

Proof. Let $v = v_1, v_2, \dots, v_n$ be a Hermite reduced basis for N . By completing squares, one obtains $f(x_1, \dots, x_n) = Q(x_1 v_1 + \dots + x_n v_n) = h_1(x_1 + c_{12}x_2 + \dots + c_{1n}x_n)^2 + g(x_2, \dots, x_n)$ where $g(x_2, \dots, x_n) = Q(x_2 \Phi_v(v_2) + \dots + x_n \Phi_v(v_n)) = h_2(x_2 + c_{23}x_3 + \dots + c_{2n}x_n)^2 + \dots + h_n x_n^2$. The quadratic forms f and g are Hermite reduced and therefore are in the Siegel domains $\mathcal{S}_n(4/3, 1/2)$ and $\mathcal{S}_{n-1}(4/3, 1/2)$ respectively. Thus the quotients $h_j/\mu_j, \mu_j/h_j, j = 1, \dots, n$, and $h_{j+1}/\mu'_j, \mu'_j/h_{j+1}, j = 1, \dots, n-1$ are bounded by a constant depending only on n , and the lemma follows (see, e.g., [1], page 266).

2. PROOF OF MAIN RESULT

We can extend T by adjoining a finite set of primes depending only on M so that for any $p \notin T$, p is odd and M_p is unimodular. We proceed with induction on n . We first make some preliminary adjustments to facilitate the proof. Choose a positive integer $e = e(M, s)$ so that the following holds at $p \in T$ ([6], Cor 5.4.3):

(*) if x and y are primitive vectors in M_p satisfying $x \equiv y \pmod{p^e M_p}$ and $Q(x) = Q(y)$ then there exists $\phi_p \in O^+(M_p)$ with $\phi_p(x) = y$ and $\phi_p \equiv 1 \pmod{p^s M_p}$.

We now define the functions $c_i(M, T, s, \lambda_{i-1})$ by induction on n . The case $n = 1$ is covered by Theorem 2.1 of [2] (for $m = 4$) and [5] (for $m \geq 5$). Note that the statement of Theorem 2.1 of [2] requires M to be integral. However, by scaling M suitably, the theorem is then applicable and we may assume $n > 1$. For $i = 1$, we define $c_1(M, T, s)$ to be the constant obtained by applying the one dimensional result to M, T, e . For $n \geq i \geq 2$, we fix $\lambda_{i-1} \in S_{i-1}(M, n)$. Then there exists a rank n \mathbf{Z} -lattice L primitively represented by the genus of M having $i - 1$ th successive minimum λ_{i-1} . Let λ_1 be the first successive minimum of L and note that since $\lambda_1 \leq \lambda_{i-1}$, there are only finitely many choices for λ_1 (although there are possibly infinitely many choices for L). If no such choice provides a λ_1 primitively represented by M , then we define $c_i(M, T, s, \lambda_{i-1})$ to be zero. For each λ_1 primitively represented by M , we pick a primitive vector $w \in M$ such that $Q(w) = \lambda_1$. Again, there are only finitely many choices for w . Set $M' = \Phi_w(M)$, $T' = T \cup \{p: \text{ord}_p(Q(w)) \neq 0\}$ and s' any integer satisfying $s' \geq s + \text{ord}_p(\gamma_w)$. Let $c_1(M', T', s')$ and $c_{i-1}(M', T', s', \lambda'_{i-2})$ be the functions given by the induction hypothesis for the case $M', T', s', n - 1$. Now for $i = 2$, we set

$$c_2(M, T, s, \lambda_1) = C \max\{c_1(M', T', s')\}$$

where C is the constant given by Lemma 1.1 and the maximum is taking over all choices of M', T', s' . For $i \geq 3$, define

$$c_i(M, T, s, \lambda_{i-1}) = C \max\{c_{i-1}(M', T', s', \lambda'_{i-2})\}.$$

Here the maximum is taking over all choices of $M', T', s', \lambda'_{i-2} \in S_{i-2}(M', n - 1)$, $\lambda'_{i-2} \leq C\lambda_{i-1}$. There are only finitely many such λ'_{i-2} since $S_{i-2}(M', n - 1) \subseteq s(M')$.

To complete the proof, let N, f_p be as given in the statement of the theorem. Let $\mu_1 \leq \dots \leq \mu_n$ be the successive minima of N . Suppose that

$\mu_i \geq c_1(M, T, s)$ and $\mu_i \geq c_i(M, T, s, \mu_{i-1})$ for $i \geq 2$. Let $v \in N$ be a vector of norm μ_1 . Then there exists a primitive vector w in M such that $Q(w) = \mu_1$ and $w \equiv f_p(v) \pmod{p^e M_p}$ for $p \in T$. We now adjust the local representation f_p . If $p \notin T$, then p is odd and M_p is unimodular. As the vectors $f_p(v)$ and w are both primitive in M_p and have the same norm, there exists $\phi_p \in O^+(M_p)$ with $\phi_p(f_p(v)) = w$. If $p \in T$, we also have $\phi_p \in O^+(M_p)$ with $\phi_p(f_p(v)) = w$ according to (*). We can now assume that $f_p(v) = w$ for all p .

We next pass to the projections $M' = \Phi_w(M)$, $N' = \Phi_v(N)$. Define $f'_p: N'_p \rightarrow M'_p$ by

$$(**) \quad f'_p(\Phi_v(u)) = \Phi_w(f_p(u)).$$

It is easy to see that f'_p is a representation. Moreover, $f'_p(N'_p)$ is primitive in M'_p . For, if $t\Phi_w(y) \in f'_p(N'_p)$ with $t \in \mathbf{Z}_p$, $y \in M_p$, then $\Phi_w(\gamma_w ty) \in f_p(N_p)$ since $\gamma_w f'_p(N'_p) = \gamma_w \Phi_w(f_p(N_p)) \subseteq f_p(N_p)$. This shows that $\gamma_w ty \in f_p(N_p)$. The primitivity of $f_p(N_p)$ implies $y \in f_p(N_p)$ and hence $\Phi_w(y) \in \Phi_w(f_p(N_p)) = f'_p(N'_p)$.

Let $\mu'_1 \leq \dots \leq \mu'_{n-1}$ be the successive minima of N' . Since $\mu_i \geq c_i$ and $C\mu_{i+1} \geq \mu'_i \geq C^{-1}\mu_{i+1}$, we have $\mu'_1 \geq c_1(M', T', s')$ and $\mu'_i \geq c_i(M', T', s', \mu'_{i-1})$ for $i = 2, \dots, n-1$. By the induction hypothesis, we can find a representation $f': N' \rightarrow M'$ satisfying conditions 1 and 2 of the theorem. Now we define $f: N \rightarrow V$ by

$$f(x) = f'(\Phi_v(x)) + \frac{(x, v)}{(v, v)} w.$$

Again, it is easy to see that f preserves the quadratic forms. Note also from (**) and the induction hypothesis that for $x \in N$ and $p \in T'$, we have $f(x) - f_p(x) = f'(\Phi_v(x)) - f'_p(\Phi_v(x)) \in \gamma_w p^s M'_p \subseteq p^s M_p$. Thus, $f(x) \in M_p$ for all $p \in T'$. It is clear that $f(x) \in M_p$ for all $p \notin T'$. This shows that $f(x) \in M$ and condition 1 of the theorem is satisfied. Since $\mathbf{Z}_p v$ splits N_p and $\mathbf{Z}_p w$ splits M_p for $p \notin T'$, the primitivity of f at these p is a consequence of the fact that $f(v) = w$ and the primitivity of $f'(N')$. It follows that $f(N)$ is primitive in M and the theorem is proved.

REFERENCES

1. J. W. S. Cassel, "Rational Quadratic Forms," Academic Press, 1978.
2. J. S. Hsia and M. Jöchner, Almost strong approximations for definite quadratic spaces, to appear in *Inventiones Math.*
3. J. S. Hsia, Y. Kitaoka, and M. Kneser, Representations of positive definite quadratic forms, *J. Reine Angew. Math.* **301** (1978), 132–141.

4. M. Jöchner, Representation of positive definite quadratic forms with congruence and primitive conditions, II, *J. Number Th.* **50** (1995), 145–153.
5. M. Jöchner and Y. Kitaoka, Representation of positive definite quadratic forms with congruence and primitive conditions *J. Number Th.* **48** (1994), 88–101.
6. Y. Kitaoka, “Arithmetic of Quadratic Forms,” Cambridge University Press, 1993.
7. O. T. O’Meara, “Introduction to Quadratic Forms,” Springer-Verlag, 1973.